



2023 South Dakota Oat Variety Trial Results Madison (Organic)

Jonathan Kleinjan | SDSU Extension Agronomist
Melanie Caffè | SDSU Oat Breeder
Kevin Kirby | Agricultural Research Manager
Shawn Hawks | Agricultural Research Manager
Nicholas Hall | Agricultural Research Manager

Cooperator: Charlie Johnson
Location: 43.862298°, -97.171856°
Soil Type: Egan-Wentworth complex, 2-6% slopes
Previous crop: soybeans
Tillage: no-till
Row spacing: 8"
Seeding Rate: 1.6 million PLS/acre
Fertilizer:
-Starter: none
-Other: none
Herbicide:
-Burndown: none
-Post: none
Fungicide: none
Date seeded: 4/27/2023
Date harvested: 8/2/2023

Note: This trial site was initiated in 2019 as part of an initiative titled “Wheat and Oat Grains for Productivity in Regenerative, Organic, and Conventional Farming Systems”, sponsored by the General Mills Foundation.



**SOUTH DAKOTA STATE
UNIVERSITY EXTENSION**

2023 South Dakota Oat Variety Trial Results Madison (Organic)

Table 1. 2023 oat variety performance trial results (average of 4 replications) at Madison, SD. Entries are sorted by overall 3-year yield. Varieties yielding in the top 1/3 of the trial are boldfaced and shaded light blue.

Variety	Height (in)	Lodging* (1-5)	Test Wt (lbs)	2021 (bu/a)	2022 (bu/a)	2023 (bu/a)	2-year (bu/a)	3-year (bu/a)
Deon	31.3	1.0	36.1	not	106.5	74.6	90.6	not
SD-Buffalo	28.3	1.0	37.2	harvested	109.6	67.7	88.7	available
MN Pearl	31.0	1.0	36.3	-	105.4	71.1	88.2	-
Hayden	30.0	1.0	36.2	-	108.0	67.6	87.8	-
Goliath	36.7	1.0	37.6	-	99.5	65.3	82.4	-
Warrior	25.0	1.0	36.1	-	100.8	55.8	78.3	-
Shelby427	29.0	1.0	38.0	-	96.2	59.9	78.0	-
Rushmore	29.0	1.0	38.0	-	95.3	60.5	77.9	-
CDC Endure	31.3	1.0	35.9	-	-	76.1	-	-
CS Camden	29.0	1.0	37.9	-	-	74.1	-	-
2018Y1315	25.7	1.0	37.5	-	-	67.5	-	-
2018Y0689	28.7	1.0	37.8	-	-	64.1	-	-
Trial Average#	30.3	1.0	37.1	-	101.7	66.9	84	-
LSD(0.05)†	-	-	2.7	-	9.8	8.3	-	-
C.V.%‡	-	-	-	-	-	7.6	-	-

* Lodging score: 1, perfectly standing; to 5, completely flat.

Trial averages may include values from experimental lines that are not reported.

† Value required (≥LSD) to determine if varieties are significantly different from one another.

‡ C.V. is a measure of variability or experimental error, 15% or less is considered acceptable.