

SOUTH DAKOTA STATE UNIVERSITY College of Agriculture, Food and Environmental Sciences

2023 South Dakota Corn Silage Trial Results Brookings

Jonathan Kleinjan | SDSU Extension Agronomist & Crop Performance Testing (CPT) Director Kevin Kirby | Agricultural Research Manager Shawn Hawks | Agricultural Research Manager

Location:	SDSU Campus - Brookings, SD 44.320784°, -96.771318°
Cooperator:	South Dakota State University Research Farm
Soil Type:	Barnes clay loam, 0-2% slope
Fertilizer:	90 lb/acre 30-10-10 starter + 134-70-50-24S preplant broadcast
Previous crop:	winter wheat
Tillage:	conventional
Row spacing:	30 inches
Seeding Rate:	33,500/acre
Herbicide:	Pre: 2.25 pt/acre Fearless
	Post: 3 oz/acre Bellum + 1 pt/acre Atrazine + qt/acre Roundup
Date seeded:	5/3/2023
Date harvested:	9/6/2023

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.

Learn more at extension.sdstate.edu.



2023 South Dakota Corn Silage Trial Results Brookings

South Dakota State University Extension

Table 1. Corn silage hybrid variety performance results (average of three replications) at Volga, SD (green chop samples).

Hybrid Informantion			Agonomic & Nutritional Performance												
Brand	Hybrid	Maturity Rating	Harvest Population ¹	Harvested ² (T/A)	DM ³ (%)	DM⁴ (T/A)	CP⁵ (%DM)	Starch ⁶ (%DM)	Lignin ⁷ (%DM)	WSC ⁸ (%DM)	NDF ⁹ (%DM)	NDFD30 ¹⁰ (%NDF)	NDFD240 ¹¹ (%NDF)	Milk2006 ¹² (lbs/T DM)	ISU Beef ¹³ (lbs/T DM)
Channel	203-83STXRIB	103	31800	28.8	47.1	10.1	7.6	40.9	2.5	4.6	36.2	57.5	71.4	3544	286
Check	DKC48-95RIB	98	33000	29.1	48.3	10.2	7.1	37.0	3.4	4.4	42.7	53.2	69.2	3232	235
Dairyland Seed	HIDF-3855Q	98	32400	29.0	47.4	10.2	7.8	36.4	3.1	4.7	40.5	52.5	69.2	3222	239
Dairyland Seed	HIDF-4545Q	102	32800	27.9	46.4	9.8	8.1	32.4	3.5	4.6	44.5	53.7	69.4	3146	223
Dekalb	DKC101-33RIB	101	30300	29.5	48.3	10.3	7.5	38.8	2.8	4.3	39.7	56.4	72.7	3372	266
Dekalb	DKC105-33RIB	105	30300	28.0	48.2	9.8	7.5	41.2	2.8	4.5	36.4	55.0	71.3	3444	277
Dekalb	DKC49-24RIB	99	32500	26.6	48.0	9.3	7.5	39.0	3.1	4.4	39.1	51.8	68.5	3281	245
Proseed	LFY 101	101	28300	26.8	43.6	9.4	8.5	29.1	3.5	4.3	46.7	53.0	69.0	3115	208
Proseed	STS 106	106	30500	31.8	39.6	11.1	8.5	33.7	3.4	5.0	41.8	50.6	67.1	3346	225
Renk Seed	RK60XPWE	94	31100	27.0	45.3	9.4	7.9	38.9	2.8	4.5	38.0	56.0	72.2	3417	267
Renk Seed	RK811PWE	104	29900	26.1	42.5	9.1	7.7	31.5	3.4	4.1	45.9	54.8	71.1	3261	224
Renk Seed	RK842VT2P	105	33300	27.7	43.1	9.7	7.6	32.7	3.5	4.2	45.0	53.9	69.5	3268	224
Renk Seed	RK895DGVT2P	105	30600	27.2	42.0	9.5	7.9	34.8	3.4	4.4	42.3	51.7	68.0	3277	222
Trial Average			31300	28.1	45.3	9.8	7.8	35.9	3.2	4.5	41.4	53.8	69.9	3302	242
LSD(0.05)†			1600	5.1	5.6	1.8	0.6	5.6	0.8	0.5	6.1	6.5	6.0	272	54
	istics are explained on p SD) to determine if varie		nificantly diffe	rent from one	another.										

SDSU Extension is an equal opportunity provider and employer in accordance with the non discrimination policies of South Dakota State University, South Dakota Board of Regents and United States Department of Agriculture. © 2023



South Dakota State University Extension

¹Plant population at harvest (plants/acre).

² Tons per acre harvested corrected to 65% moisture.

³ Dry matter (DM) percentage of harvested corn silage.

⁴ Tons per acre of dry matter (DM).

⁵ Crude protein (CP), % of dry matter.

⁶ Starch, % of dry matter.

⁷ Lignin, % of dry matter.

⁸ Water Soluble Carbohydrates (WSC), % of dry matter.

⁹Neutral detergent fiber (NDF), % of dry matter.

¹⁰ 30 hour digestibility of NDF (NDFD30) is the amount of NDF digested in 30 hours as a percentage of NDF.

¹¹ 240 hour digestibility of NDF (NDFD240) is the amount of NDF digested in 240 hours as a percentage of NDF.

¹² Milk2006 is the prediction of the amount of milk produced per ton of corn silage dry matter.

¹³ ISU Beef is the prediction of the amount of beef produced per ton of corn silage dry matter.

Procedure:

Corn was harvested for silage by hand cutting at 6 – 8 inches from the ground. Material was weighed. Material was chopped through a chipper/shredder. Green chop samples were frozen. Samples submitted to a commercial laboratory for nutrient analyses using calibrated NIR instrumentation.

For Further Information:

Dr. Jonathan Kleinjan, SDSU Extension Agronomist (605) 688-4211 jonathan.kleinjan@sdstate.edu

2023 South Dakota Corn Silage Trial Results Brookings